Appendix C - Aquatic Invasive Species of Concern

Non-native, Aquatic Invasive Plant Species

EURASIAN WATERMILFOIL(EWM)

EWM is a submersed aquatic plant native to Europe, Asia, and northern Africa (Figure 1). It is the only non-native milfoil in Wisconsin. Like the native milfoils, the Eurasian variety has slender stems whorled by submersed feathery leaves and tiny flowers produced above the water surface. The flowers are located in the axils of the floral bracts, and are either four-petaled or without petals. The leaves are threadlike, typically uniform in diameter, and aggregated into a submersed terminal spike. The stem thickens below the inflorescence and doubles its width further down, often curving to lie parallel with the water surface. The fruits are four-jointed nut-like bodies. Without flowers or fruits, EWM is difficult to distinguish from Northern water milfoil. EWM has 9-21 pairs of leaflets per leaf, while Northern milfoil typically has 7-11 pairs of leaflets. Coontail is often mistaken for the milfoils, but does not have individual leaflets.

EWM grows best in fertile, fine-textured, inorganic sediments. In less productive lakes, it is restricted to areas of nutrient-rich sediments. It has a history of becoming dominant in eutrophic, nutrient-rich lakes, although this pattern is not universal. It is an opportunistic species that prefers highly disturbed lake beds, lakes receiving nitrogen and phosphorous-laden runoff, and heavily used lakes. Optimal growth occurs in alkaline systems with a high concentration of dissolved inorganic carbon. High water temperatures promote multiple periods of flowering and fragmentation.

Unlike many other plants, EWM does not rely on seed for reproduction. Its seeds germinate poorly under natural conditions. It reproduces by fragmentation, allowing it to disperse over long distances. The plant produces fragments after fruiting once or twice during the summer. These shoots may then be carried downstream by water currents or inadvertently picked up by boaters. EWM is readily dispersed by boats, motors, trailers, bilges, live wells, and bait buckets; and can stay alive for weeks if kept moist.

Once established in an aquatic community, milfoil reproduces from shoot fragments and stolons (runners that creep along the lake bed). As an opportunistic species, EWM is adapted for rapid growth early in spring. Stolons, lower stems, and roots persist over winter and store the carbohydrates that help milfoil claim the water column early in spring, photosynthesize, divide, and form a dense leaf canopy that shades out native aquatic plants. Its ability to spread rapidly by fragmentation and effectively block out sunlight needed for native plant growth often results in monotypic stands. Monotypic stands of EWM provide only a single habitat, and threaten the integrity of aquatic communities in a number of ways; for example, dense stands disrupt predator-prey relationships by fencing out larger fish, and reducing the number of nutrient-rich native plants available for waterfowl.

Dense stands of EWM also inhibit recreational uses like swimming, boating, and fishing. Some stands have been dense enough to obstruct industrial and power generation water intakes. The visual impact that greets the lake user on milfoil-dominated lakes is the flat yellow-green of matted vegetation, often prompting the perception that the lake is "infested" or "dead". Cycling of nutrients from sediments to the water column by EWM may lead to deteriorating water quality and algae blooms in infested lakes.

Figure 1: Eurasian watermilfoil

CURLY-LEAF PONDWEED

CLP is an invasive aquatic perennial that is native to Eurasia, Africa, and Australia (Figure 2). It was accidentally introduced to United States waters in the mid-1880s by hobbyists who used it as an aquarium plant. The leaves are reddish-green, oblong, and about 3 inches long, with distinct wavy edges that are finely toothed. The stem of the plant is flat, reddish-brown and grows from 1 to 3 feet long. CLP is commonly found in alkaline and high nutrient waters, preferring soft substrate and shallow water depths. It tolerates low light and low water temperatures. It has been reported in all of the lower 48 states.

CLP spreads through burr-like winter buds (turions) (Figure 2), which are moved among waterways. These plants can also reproduce by seed, but this plays a relatively small role compared to the vegetative reproduction through turions. New plants form under the ice in winter, making curly-leaf pondweed one of the first nuisance aquatic plants to emerge in the spring. It becomes invasive in some areas because of its tolerance for low light and low water temperatures. These tolerances allow it to get a head start on and out-compete native plants in the spring. In mid-summer, when most aquatic plants are growing, CLP plants are dying off. Plant die-offs may result in a critical loss of dissolved oxygen. Furthermore, the decaying plants can increase nutrients which contribute to algal blooms, as well as create unpleasant stinking messes on beaches. CLP forms surface mats that interfere with aquatic recreation.

Figure 2: Curly-leaf pondweed and its turions

PURPLE LOOSESTRIFE

Purple loosestrife (Figure 3) is a perennial herb 3-7 feet tall with a dense bushy growth of 1-50 stems. The stems, which range from green to purple, die back each year. Showy flowers that vary from purple to magenta possess 5-6 petals aggregated into numerous long spikes, and bloom from August to September. Leaves are opposite, nearly linear, and attached to four-sided stems without stalks. It has a large, woody taproot with fibrous rhizomes that form a dense mat. By law, purple loosestrife is a nuisance species in Wisconsin. It is illegal to sell, distribute, or cultivate the plants or seeds, including any of its cultivars.

Purple loosestrife is a wetland herb that was introduced as a garden perennial from Europe during the 1800's. It is still promoted by some horticulturists for its beauty as a landscape plant, and by beekeepers for its nectar-producing capability. Currently, more than 20 states, including Wisconsin have laws prohibiting its importation or distribution because of its aggressively invasive characteristics. It has since extended its range to include most temperate parts of the United States and Canada. The plant's reproductive success across North America can be attributed to its wide tolerance of physical and chemical conditions characteristic of disturbed habitats, and its ability to reproduce prolifically by both seed dispersal and vegetative propagation. The absence of natural predators, like European species of herbivorous beetles that feed on the plant's roots and leaves, also contributes to its proliferation in North America.

Purple loosestrife was first detected in Wisconsin in the early 1930's, but remained uncommon until the 1970's. It is now widely dispersed in the state, and has been recorded in 70 of Wisconsin's 72 counties. Low densities in most areas of the state suggest that the plant is still in the pioneering stage of establishment. Areas of heaviest infestation are sections of the Wisconsin River, the extreme southeastern part of the state, and the Wolf and Fox River drainage systems.

This plant's optimal habitat includes marshes, stream margins, alluvial flood plains, sedge meadows, and wet prairies. It is tolerant of moist soil and shallow water sites such as pastures and meadows, although established plants can tolerate drier conditions. Purple loosestrife has also been planted in lawns and gardens, which is often how it has been introduced to many of our wetlands, lakes, and rivers.

Purple loosestrife can germinate successfully on substrates with a wide range of pH. Optimum substrates for growth are moist soils of neutral to slightly acidic pH, but it can exist in a wide range of soil types. Most seedling establishment occurs in late spring and early summer when temperatures are high.

Purple loosestrife spreads mainly by seed, but it can also spread vegetatively from root or stem segments. A single stalk can produce from 100,000 to 300,000 seeds per year. Seed survival is up to 60-70%, resulting in an extensive seed bank. Mature plants with up to 50 shoots grow over 2 meters high and produce more than two million seeds a year. Germination is restricted to open, wet soils and requires high temperatures, but seeds remain viable in the soil for many years. Even seeds submerged in water can live for approximately 20 months. Most of the seeds fall near the parent plant, but water, animals, boats, and humans can transport the seeds long distances. Vegetative spread through local perturbation is also characteristic of loosestrife; clipped, trampled, or buried stems of established plants may produce shoots and roots. Plants may be quite large and several years old before they begin flowering. It is often very difficult to locate non-flowering plants, so monitoring for new invasions should be done at the beginning of the flowering period in mid-summer.

Any sunny or partly shaded wetland is susceptible to purple loosestrife invasion. Vegetative disturbances such as water drawdown or exposed soil accelerate the process by providing ideal conditions for seed germination. Invasion usually begins with a few pioneering plants that build up a large seed bank in the soil for several years. When the right disturbance occurs, loosestrife can spread rapidly, eventually taking over the entire wetland. The plant can also make morphological adjustments to accommodate changes in the immediate environment; for example, a decrease in light level will trigger a change in leaf morphology. The plant's ability

to adjust to a wide range of environmental conditions gives it a competitive advantage; coupled with its reproductive strategy, purple loosestrife tends to create monotypic stands that reduce biotic diversity. Purple loosestrife displaces native wetland vegetation and degrades wildlife habitat. As native vegetation is displaced, rare plants are often the first species to disappear. Eventually, purple loosestrife can overrun wetlands thousands of acres in size, and almost entirely eliminate the open water habitat. The plant can also be detrimental to recreation by choking waterways.

Figure 3: Purple loosestrife

REED CANARY GRASS

Reed canary grass (Figure 4) is a large, coarse grass that reaches 2 to 9 feet in height. It has an erect, hairless stem with gradually tapering leaf blades 3 1/2 to 10 inches long and 1/4 to 3/4 inch in width. Blades are flat and have a rough texture on both surfaces. The lead ligule is membranous and long. The compact panicles are erect or slightly spreading (depending on the plant's reproductive stage), and range from 3 to 16 inches long with branches 2 to 12 inches in length. Single flowers occur in dense clusters in May to mid-June. They are green to purple at first and change to beige over time. This grass is one of the first to sprout in spring, and forms a thick rhizome system that dominates the subsurface soil. Seeds are shiny brown in color.

Both Eurasian and native ecotypes of reed canary grass are thought to exist in the U.S. The Eurasian variety is considered more aggressive, but no reliable method exists to tell the ecotypes apart. It is believed that the vast majority of our reed canary grass is derived from the Eurasian ecotype. Agricultural cultivars of the grass are widely planted.

Reed canary grass also resembles non-native orchard grass (*Dactylis glomerata*), but can be distinguished by its wider blades, narrower, more pointed inflorescence, and the lack of hairs on glumes and lemmas (the spikelet scales). Additionally, bluejoint grass (*Calamagrostis canadensis*) may be mistaken for reed canary in areas where orchard grass is rare, especially in the spring. The highly transparent ligule on reed canary grass is helpful in distinguishing it from the others. Ensure positive identification before attempting control.

Figure 4: Reed canary grass

HYBRID CATTAIL

Hybrid cattail (Figure 5) is native to southern but not northern Wisconsin, Narrow-leaved cattail (*Typha angustifolia*) and its hybrids with Broad-leaved cattail are becoming increasingly common in northern Wisconsin where they also tend to be invasive. Besides having narrower leaves, the exotics can be told from native cattails by having a relatively narrower and longer "hotdog-shaped" tan female cattail flower whereas the native species tends to produce a fatter and shorter "bratwurst-shaped" dark chocolate colored female flower. Narrow-leaved cattail and its hybrids also have a male flower that is separated from the female flower by a thin green stem while the native Broad-leaved cattail has its male and female flowers connected (Figure 5).


Figure 5: Hybrid cattail

YELLOW FLAG IRIS

Yellow flag iris (Figure 6) is a showy perennial plant that can grow in a range of conditions from drier upland sites, to wetlands, to floating aquatic mats. A native plant of Eurasia, it can be an invasive garden escapee in Wisconsin's natural environments. Yellow flag iris can produce many seeds that can float from the parent plant, or plants can spread vegetatively via rhizome fragments. Once established it forms dense clumps or floating mats that can alter wildlife habitat and species diversity. All parts of this plant are poisonous, which results in lowered wildlife food sources in areas where it dominates. This species has the ability to escape water gardens and ponds and grow in undisturbed and natural environments. It can grow in wetlands, forests, bogs, swamps, marshes, lakes, streams and ponds. Dense areas of this plant may alter hydrology by trapping sediment and/or blocking flow.

Yellow iris has broad, sword-shaped leaves that grow upright, tall and stiff. They are green with a slight blue-grey tint and are very difficult to distinguish from other ornamental or native iris species. Flowers are produced on a stem that can grow 3-4 feet tall among leaves that are usually as tall or taller. The flowers are showy and variable in color from almost white to a vibrant dark yellow. Flowers are between 3-4 inches wide and bloom from April to June. Three upright petals are less showy than the larger three downward pointing sepals, which may have brown to purple colored streaks. Seeds are produced in fruits that are 6-angled capsules, 2-4 inches long. Each fruit may have over 100 seeds that start pale before turning dark brown. Each seed has a hard outer casing with a small air space underneath, which allows the seeds to float. The roots are thick, fleshy pink-colored rhizomes spread extensively in good conditions, forming thick mats that can float on the surface of the water.

When not flowering, yellow flag iris could be easily confused with the native blue flag iris (Iris versicolor) as well as other ornamental irises that are not invasive. Blue flag iris is usually smaller and does not tend to form as dense clumps or floating mats. When not flowering or showing fruiting bodies, yellow flag iris may be confused with other wetland plants such as cattails (Typha spp.) or sweet flag (Acorus spp.) species.
Small populations may be successfully removed using physical methods. Care should be taken if hand-pulling plants as some people show skin sensitivity to plant sap and tissues. All parts of the plant should be dug out – particularly rhizomes and disposed of in a landfill or by burning. Cutting the seed heads may help decrease the plant spreading.

Aquatic formulas of herbicides may be used to control yellow flag iris, however, permits may be needed. Foliar spray, cut stem/leaf application and hand swiping of herbicide have all shown effectiveness.

Figure 6: Yellow flag iris

Non-native Aquatic Invasive Animal Species

CHINESE MYSTERY SNAILS

The Chinese mystery snails and the banded mystery snails (Figure 7) are non-native snails that have been found in a number of Wisconsin lakes. There is not a lot yet known about these species, however, it appears that they have a negative effect on native snail populations. The mystery snail's large size and hard operculum (a trap door cover which protects the soft flesh inside), and their thick hard shell make them less edible by predators.

The female mystery snail gives birth to live crawling young. This may be an important factor in their spread as it only takes one impregnated snail to start a new population. Mystery snails thrive in silt and mud areas although they can be found in lesser numbers in areas with sand or rock substrates. They are found in lakes, ponds, irrigation ditches, and slower portions of streams and rivers. They are tolerant of pollution and often thrive in stagnant water areas. Mystery snails can be found in water depths of 0.5 to 5 meters (1.5 to 15 feet). They tend to reach their maximum population densities around 1-2 meters (3-6 feet) of water depth. Mystery snails do not eat plants. Instead, they feed on detritus and in lesser amounts on algae and phytoplankton. Thus removal of plants along the shoreline area will not reduce the abundance of mystery snails.

Lakes with high densities of mystery snails often see large die-offs of the snails. These die-offs are related to the lake's warming coupled with low oxygen (related to algal blooms). Mystery snails cannot tolerate low oxygen levels. High temperatures by themselves seem insufficient to kill the snails as the snails could move into deeper water.

Many lake residents are worried about mystery snails being carriers of the swimmer's itch parasite. In theory they are potential carriers, however, because they are an introduced species and did not evolve as part of the lake ecosystem, they are less likely to harbor the swimmer's itch parasites.

Figure 7: Chinese Mystery Snails

ZEBRA MUSSELS

Zebra mussels (Figure 8) are an invasive species that have inhabited Wisconsin waters and are displacing native species, disrupting ecosystems, and affecting citizens' livelihoods and quality of life. They hamper boating, swimming, fishing, hunting, hiking, and other recreation, and take an economic toll on commercial, agricultural, forestry, and aquacultural resources. The zebra mussel is a tiny (1/8-inch to 2-inch) bottom-dwelling clam native to Europe and Asia. Zebra mussels were introduced into the Great Lakes in 1985 or 1986, and have been spreading throughout them since that time. They were most likely brought to North America as larvae in ballast water of ships that traveled from fresh-water Eurasian ports to the Great Lakes. Zebra mussels look like small clams with a yellowish or brownish D-shaped shell, usually with alternating dark- and light-colored stripes. They can be up to two inches long, but most are under an inch. Zebra mussels usually grow in clusters containing numerous individuals.

Zebra mussels feed by drawing water into their bodies and filtering out most of the suspended microscopic plants, animals and debris for food. This process can lead to increased water clarity and a depleted food supply for other aquatic organisms, including fish. The higher light penetration fosters growth of rooted aquatic plants which, although creating more habitat for small fish, may inhibit the larger, predatory fish from finding their food. This thicker plant growth can also interfere with boaters, anglers and swimmers. Zebra mussel infestations may also promote the growth of blue-green algae, since they avoid consuming this type of algae but not others.

Zebra mussels attach to the shells of native mussels in great masses, effectively smothering them. A survey by the Army Corps of Engineers in the East Channel of the Mississippi River at Prairie du Chien revealed a substantial reduction in the diversity and density of native mussels due to Zebra Mussel infestations. The East Channel provides habitat for one of the best mussel beds in the Upper Mississippi River. Future efforts are being considered to relocate such native mussel beds to waters that are less likely to be impacted by zebra mussels.

Once zebra mussels are established in a water body, very little can be done to control them. It is therefore crucial to take all possible measures to prevent their introduction in the first place. Some of the preventative and physical control measures include physical removal, industrial vacuums, and back flushing.

Chemical applications include solutions of chlorine, bromine, potassium permanganate and even oxygen deprivation. An ozonation process is under investigation (patented by Bollyky Associates Inc.) which involves the pumping of high concentrations of dissolved ozone into the intake of raw water pipes. This method only works in controlling veligers, and supposedly has little negative impacts on the ecosystem. Further research on effective industrial control measures that minimize negative impacts on ecosystem health is needed.

In the fall of 2016, zebra mussels were found in Big and Middle McKenzie Lakes on the Washburn/Burnett County line southwest of Gilmore Lake. Since then, zebra mussels have also been identified in Deer Lake, Balsam Lake, and Wapogasset Lake in Polk County and in Bass Lake in St, Croix County. They are also in Lake Superior.

Figure 8: Zebra Mussels

RUSTY CRAYFISH

Rusty crayfish (Figure 9) live in lakes, ponds and streams, preferring areas with rocks, logs and other debris in water bodies with clay, silt, sand or rocky bottoms. They typically inhabit permanent pools and fast moving streams of fresh, nutrient-rich water. Adults reach a maximum length of 4 inches. Males are larger than females upon maturity and both sexes have larger, heartier, claws than most native crayfish. Dark "rusty" spots are usually apparent on either side of the carapace, but are not always present in all populations. Claws are generally smooth, with grayish-green to reddish-brown coloration. Adults are opportunistic feeders, feeding upon aquatic plants, benthic invertebrates, detritus, juvenile fish and fish eggs.

The native range of the rusty crayfish includes Ohio, Tennessee, Kentucky, Indiana, Illinois and the entire Ohio River basin. However, this species may now be found in Michigan, Massachusetts, Missouri, Iowa, Minnesota, New York, New Jersey, Pennsylvania, Wisconsin, New Mexico and the entire New England state area (except Rhode Island). The Rusty crayfish has been a reported invader since at least the 1930's. Its further spread is of great concern since the prior areas of invasion have led to severe impacts on native flora and fauna. It is thought to have spread by means of released game fish bait and/or from aquarium release. Rusty crayfish are also raised for commercial and biological harvest.

Rusty crayfish reduce the amount and types of aquatic plants, invertebrate populations, and some fish populations--especially bluegill, smallmouth and largemouth bass, lake trout and walleye. They deprive native fish of their prey and cover and out-compete native crayfish. Rusty crayfish will also attack the feet of swimmers. On the positive side, rusty crayfish can be a food source for larger game fish and are commercially harvested for human consumption.

Rusty crayfish may be controlled by restoring predators like bass and sunfish populations. Preventing further introduction is important and may be accomplished by educating anglers, trappers, bait dealers and science teachers of their hazards. Use of chemical pesticides is an option, but does not target this species and will kill other aquatic organisms.

It is illegal to possess both live crayfish and angling equipment simultaneously on any inland Wisconsin water (except the Mississippi River). It is also illegal to release crayfish into a water of the state without a permit.



Figure 9: Rusty Crayfish and identifying characteristics